

# Cigarette smoking and the onset and persistence of depression among adults in the United States: 1994–2005

Jafar Bakhshaie<sup>a</sup>, Michael J. Zvolensky<sup>a,b</sup>, Renee D. Goodwin<sup>c,d,\*</sup>

<sup>a</sup>Department of Psychology, University of Houston, 126 Heyne Building, Houston, TX, 77204, United States

<sup>b</sup>Department of Behavioral Sciences, University of Texas MD Anderson Cancer Center, Houston, TX, United States

<sup>c</sup>Department of Psychology, Queens College and The Graduate Center, City University of New York (CUNY), Queens, NY, United States

<sup>d</sup>Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, United States

## Abstract

**Background:** The present study investigated the relationship between daily cigarette smoking and risk of onset and persistence of major depressive disorder (MDD) over a 10-year period among adults in the United States and whether successful smoking cessation reduced the risk for MDD.

**Method:** Data were drawn from the Midlife Development in the United States Survey (MIDUS;  $n = 2101$ ) Waves I and II. Logistic regressions were used to investigate the relations between smoking and the onset and persistence of MDD, adjusting for demographic characteristics and substance use problems.

**Results:** Daily smoking in 1994 [OR = 1.9 (1.2–3.2)] and persistent daily smoking (in 1994 and 2005) [OR = 2.2 (1.3–3.7)] were both associated with a significantly increased likelihood of MDD in 2005. Additionally, abstinence, compared to daily smoking, for more than 10 years significantly reduced the risk of MDD in 2005 [OR = 0.5 (0.3–0.87)] and persistent MDD in 1994 and 2005; [OR = 0.5 (0.3–0.87)].

**Conclusions:** Findings from this study provide new insights into the role of smoking in the onset and persistence of MDD. Namely, among those in mid-adulthood, smoking is associated with greater MDD risk and quitting may help to reduce such risk. These results suggest that there may be merit in targeting smoking to reduce the risk of MDD and the mental health benefits of quitting smoking in the form of reduced risk of MDD could usefully be added to common information listed as reasons to quit.

© 2015 Published by Elsevier Inc.

## 1. Introduction

There is substantial comorbidity between major depressive disorder (MDD) and smoking [1]. For example, Lasser and colleagues (2000) reported that approximately 60% of those with a lifetime history of MDD are current or past smokers [2]. Conversely, smokers relative to non-smokers have significantly higher rates of MDD [3]. Other work suggests that smokers who are nicotine dependent are significantly more likely to have a history of MDD [4].

There are a variety of streams of empirical work that suggest smoking–MDD relations are bi-directional. First, MDD may influence the developmental progression to smoking via self-medication or mood management mechanisms. For

example, young adults with a history of MDD compared to those without such a history are significantly more likely to begin smoking [5]. Yet, other studies have not found such MDD to smoking onset relations [6]. Second, some work suggest smoking influences the onset and course of MDD; a pathway theorized to be influenced by neuro-physiologic processes altered by smoking [7,8]. In fact, daily smoking is associated with increased risk for MDD in longitudinal studies [9–11]. Finally, there is mixed evidence that a history of MDD is related to lower odds of cessation [12–14]. Yet, moderate to severe levels of depressive symptoms prior to smoking cessation treatment are significantly associated with poorer cessation outcomes [15–17]. In addition, therapeutic approaches that include depression treatment components compared to standard smoking cessation protocols appear to improve odds of quitting [18–20].

Despite progress in better understanding smoking–MDD developmental relations, there are at least three key limitations to past work. First, in terms of the developmental path from cigarette smoking to MDD, longer-term outcomes

\* Corresponding author at: Department of Psychology, Queens College and The Graduate Center, City University of New York (CUNY), 65–30 Kissena Boulevard, Queens, New York 11367. Tel.: +1 718 997 3247; fax: +1 212 342 5168.

E-mail address: [renee.goodwin@qc.cuny.edu](mailto:renee.goodwin@qc.cuny.edu) (R.D. Goodwin).

have been infrequently modeled. Indeed, most studies on the impact of smoking on the risk of future MDD have employed follow-up periods that range only from 6 months to 5 years [5,21,22]. Thus, there is a need for the replication and extension of past work using longer periods of time. Second, past studies have not yet explored the explanatory utility of persistence of smoking (i.e., ‘stable’ smoking measured between at least two time points) over time in relation to MDD. Such information is practically and theoretically important, as it will help shed novel insight into the role of persistent smoking behavior and relative risk for new and persistent cases of MDD. Finally, past work has infrequently examined how smoking cessation relates to the risk of recent incidence and persistence of MDD. Thus, the clinical significance of smoking as a risk factor for MDD prevention/intervention has not been fully explicated. Finally, a majority of past work has been completed in younger persons [23], and therefore, relatively little is known about these relations over the course of adulthood. This limitation is important because smoking is the leading cause of premature death among older adults [24,25].

Together, the current investigation examined the relation between smoking and MDD over a 10 year period. First, it was hypothesized that after controlling for gender [26,27] socioeconomic status [28–30], and substance use problems [31,32], daily smoking (compared to non-smoking) would be associated with increased likelihood of MDD. Second, it was hypothesized that successful smoking cessation (compared to persistent smoking) would be associated with decreased likelihood of MDD.

## 2. Methods

### 2.1. Participants

Data were drawn from the two waves of the Midlife Development in the United States Survey (MIDUS) [33,34]. The MacArthur Midlife Research Network collected wave I data from 1994 to 1995 and wave II data from 2004 to 2006; a national survey of Americans in adulthood that investigated behavioral, psychological, and social factors related to physical and mental health. Wave I consisted of a nationally representative multistage probability sample (main sample) of community-dwelling English speakers in the continental United States ( $n = 3032$ ). Participants who completed the telephone interview were mailed a self-administered questionnaire. The response rate from the mailed questionnaire was 86.6%, yielding an overall response rate of 61% ( $.70 * .87 = .61$ ). Approximately 70% of wave I participants took part in the wave II survey collected by the Institute on Aging at the University of Wisconsin–Madison and supported by the National Institute on Aging (2004–2006). Wave II participants completed a 30-min telephone interview and a self-administered questionnaire was mailed to them. Of the 3032 participants from wave I, 2101 completed the wave II telephone surveys (response rate of 69.5%). For

this study, we analyzed only data from those who participated in the wave I main sample who completed both the phone and mail-in surveys, participated in the wave II survey, and had complete information for wave II outcome variables.

### 2.2. Measures

#### 2.2.1. Demographic characteristics

At wave 1, basic socio-demographic characteristics were obtained using a self-report questionnaire. These included age, race, gender, marital status and level of formal education obtained.

#### 2.2.2. Smoking

At each wave, participants who reported ever smoking at least one cigarette were asked whether they had ever smoked regularly (“regularly—meaning at least a few cigarettes every day”), the age they were when they last smoked regularly and whether they currently smoke regularly (“Do you smoke cigarettes regularly now?”). Those who reported no lifetime smoking were considered *lifetime non-smoker*. Those who reported lifetime daily smoking and were not current daily smokers at wave 1 or 2 were considered *former daily smokers* with successful abstinence for more than 10 years. Those who reported current daily smoking at waves 1 and 2 were considered *persistent smokers*.

#### 2.2.3. Major depression

Psychiatric diagnoses of MIDUS were based on the Composite International Diagnostic Interview Short Form scales (CIDI-SF; 38). CIDI-SF is a series of diagnostic-specific scales that were developed from item level analyses of the Composite International Diagnostic Interview questions during the National Comorbidity Survey [35]. CIDI-SF scales were designed to reproduce the full CIDI diagnoses as exactly as possible, with only a small subset of the original questions. Validity data suggest strong relationship between CIDI-SF-based and the full CIDI-based diagnoses [36]. These measures were used in waves 1 and 2 for the presence of the disorders during the past-12 month. Recent incidence of MDD by 2005 was defined as having past 12-month disorder at wave 2 but not at wave 1. Persistent MDD was defined as having past 12-month disorder at both wave 1 and wave 2.

### 2.3. Statistical analysis

Differences in demographic characters between lifetime non-smokers, former daily smokers (abstinence for more than 10 years after history of daily smoking), and persistent smokers were identified using chi-square analyses. Logistic regression analyses were utilized to evaluate the relationships between (a) daily smokers (1994) vs. lifetime non-smokers; (b) persistent smokers vs. lifetime non-smokers; (c) former daily smokers (successful quitters) vs. lifetime non-smokers; (d) former daily smokers (successful quitters) vs. persistent smokers, and odds of (1) past 12-month MDD by 2005; (2) persistent/recurring MDD in

Table 1

Demographic characteristics<sup>a</sup> associated with remitted/former daily smoking (i.e., abstinence for more than 10 years) and persistent smoking among adults in the community.

	Lifetime non- smokers (n = 518)	Former daily smokers "10+ years abstinence" (n = 532)	p-Value	Persistent smokers (n = 356)	p-Value
Age-mean (SD)	43.8 (13.5)	49.9 (12.3)	<0.0001	41.3 (10.5)	0.003
Gender					
Male	(177) 32.10%	(290) 54.50%	<0.0001	(176) 49.40%	<0.0001
Female	(342) 65.90%	(242) 45.50%		(180) 50.60%	
Race			0.1		0.4
White	(416) 87.40%	(451) 91.10%		(283) 90.10%	
Black	(32) 6.70%	(29) 5.90%		(21) 6.70%	
Native American	(3) 0.60%	(1) 0.20%		(1) 0.30%	
Asian or Pacific Islander	(9) 1.90%	(2) 0.40%		(1) 0.30%	
Other	(12) 2.50%	(7) 1.40%		(7) 2.20%	
Multiracial	(4) 0.80%	(5) 1.00%		(1) 0.30%	
Education			<0.0001		<0.0001
Some grade school	(52) 10.00%	(77) 14.50%		(77) 21.60%	
High school graduate	(141) 27.20%	(172) 32.30%		(149) 41.90%	
Some college	(141) 18.70%	(149) 28.00%		(98) 27.50%	
Bachelor's degree	(97) 18.70%	(86) 16.20%		(20) 5.60%	
Some graduate school or beyond	(88) 18.70%	(86) 16.20%		(12) 3.40%	
Marital status			0.026		<0.0001
Married	(361) 69.80%	(385) 72.50%		(227) 63.60%	
Separated	(7) 1.40%	(8) 1.50%		(10) 2.80%	
Divorced	(54) 10.40%	(69) 13.00%		(75) 21.00%	
Widowed	(46) 8.90%	(45) 8.50%		(17) 4.80%	
Never married	(49) 9.50%	(24) 4.50%		(28) 7.80%	

<sup>a</sup> Demographic characteristics reported at wave 1.

1994 and 2005. Analyses were adjusted for socio-demographic factors and alcohol and drug use problems.

### 3. Results

#### 3.1. Demographic characteristics and smoking

Compared with non-smokers, persistent smokers were less likely to be female, have lower age or fewer years of education, less likely to be married, and more likely to be separated, divorced, or never married (see Table 1). More-

over, compared to non-smokers, smokers who successfully quit were less likely to be female, had less education, and were more likely to be divorced and be older (see Table 1).

#### 3.2. Daily smoking and MDD

Smoking at 1994 compared to non-smoking was associated with an increased risk of incidence of MDD by 2005 and persistence of MDD from 1994 to 2005 (see Table 2). These associations remained significant after adjusting for the covariates.

Table 2  
Current daily smoking in 1994, and persistent daily smoking (in 1994 and 2005) and risk of depression onset and persistence in 2005, both compared to lifetime non-smoking.

	Lifetime non-smokers ( <i>n</i> = 518)	Current daily smokers 1994 ( <i>n</i> = 525)	OR (95% CI)	AOR <sup>a</sup> (95% CI)	AOR <sup>b</sup> (95% CI)	Persistent daily smokers <sup>c</sup> ( <i>n</i> = 356)	OR (95% CI)	AOR <sup>a</sup> (95% CI)	AOR <sup>b</sup> (95% CI)
<b>MDD 2005 Only</b>	6.1% (28)	11.2% (46)	<b>1.9 (1.2, 3.2)</b>	<b>2.2 (1.4, 3.7)</b>	<b>1.8 (1.1, 3.1)</b>	13.5% (38)	<b>2.4 (1.4, 4.0)</b>	<b>2.8 (1.6, 4.7)</b>	<b>2.3 (1.3, 4.0)</b>
<b>MDD 1994 + 2005</b>	5.1% (23)	10.4% (42)	<b>2.2 (1.3, 3.7)</b>	<b>2.6 (1.5, 4.5)</b>	<b>2.5 (1.4, 4.3)</b>	10.9% (30)	<b>2.3 (1.3, 4.0)</b>	<b>2.7 (1.5, 4.9)</b>	<b>2.6 (1.4, 4.8)</b>

Bold = *p* < 0.05.

<sup>a</sup> AOR=age and gender.

<sup>b</sup> AOR=age, gender and alcohol/drug use problems at wave 1 (1994).

<sup>c</sup> Smoking at both waves (1994 and 2005).

Table 3  
Former daily smoking (i.e., abstinence for more than 10 years) and risk of depression onset and persistence in 2005, compared to persistent daily smoking (1994 and 2005) and lifetime non-smoking.

	Former daily smokers “10+ years abstinence” ( <i>n</i> = 532)	Persistent daily smokers ( <i>n</i> = 356)	OR (95% CI)	AOR <sup>a</sup> (95% CI)	AOR <sup>b</sup> (95% CI)	Lifetime non-smokers ( <i>n</i> = 518)	OR (95% CI)	AOR <sup>a</sup> (95% CI)	AOR <sup>b</sup> (95% CI)
<b>MDD 2005 Only</b>	7.7% (36)	13.5% (38)	<b>0.5 (0.3, 0.87)</b>	0.6 (0.4, 1.06)	0.7 (0.4, 1.2)	6.1% (28)	1.3 (0.8, 2.1)	<b>1.8 (1.03, 3.0)</b>	1.7 (0.98, 2.9)
<b>MDD 1994 + 2005</b>	4.2% (19)	10.9% (30)	<b>0.36 (0.2, 0.65)</b>	<b>0.50 (0.26, 0.93)</b>	<b>0.5 (0.26, 0.96)</b>	5.1% (23)	0.8 (0.4, 1.5)	1.2 (0.6, 2.3)	1.1 (0.6, 2.2)

Bold = *p* < 0.05.

<sup>a</sup> AOR=age and gender.

<sup>b</sup> AOR=age, gender and alcohol/drug use problems at wave 1 (1994).

### 3.3. Persistent smoking and MDD

Persistent smoking compared to non-smoking was associated with increased risk of MDD (see Table 2); these effects were significant after adjusting for the covariates.

### 3.4. Former daily smoking (sustained abstinence for more than 10 years) and MDD

Compared to persistent smoking sustained abstinence for more than 10 years was significantly related to lower risk for MDD (see Table 3). Additionally, compared to lifetime non-smokers (see Table 3), sustained abstinence for more than 10 years was not significantly associated with MDD, even after adjusting for demographic factors, and alcohol/drug use problems. The only exception to this pattern of results was a slightly higher incidence of MDD at 2005 for those who had quit compared to non-smokers, after adjusting for age and gender (see Table 3).

## 4. Discussion

Smoking is associated with significantly increased risk of the onset and persistence of MDD even after controlling for socio-demographic characteristics and substance use problems. These observations are in line with previous findings and extend them to a 10 year period among a nationally representative sample of adults in the U.S. [5,37]. Moreover, smokers who quit and remained abstinent for more than 10 years, compared to persistent smokers, demonstrated lower odds of MDD [21,38–40]. However, an alternative explanation could be that this successful quitting may have been as a result of the prior remission of MDD. Nonetheless, with replication in future work, this finding suggests quitting smoking may result in lower risk of MDD whether from an onset or maintenance perspective. This is a potentially clinically important observation that suggests there is merit to the utilization of smoking cessation as a method to prevent or reduce risk for MDD [41–43].

It is noteworthy that the risk of MDD for those who quit was not different from non-smokers. Regardless of possible explanations for this observation (i.e., whether quitting reduced the risk of MDD, or clinical improvement in MDD contributed to cessation), the extant literature has yielded inconsistent findings regarding the MDD-related prognoses of former smokers compared to non-smokers [22,27,44–53]. These inconsistencies could be due to utilization of different methodologies [49] and between-sample differences in variables like gender [27,49], comorbid anxiety [44], and genetic predisposition [54–57]. However, *even* among the studies that report higher rates of MDD for former smokers compared to lifetime non-smokers, these rates are still lower than for current daily/persistent smokers [22,27,44].

The present study has a number of limitations. First, despite utilization of longitudinal methodology, it is not possible to discern the exact chronology of some theoretic-

cally and clinically-relevant events (e.g., quitting dates, dates of onset of past/recent MDD). A useful next step in this line of work therefore would be to isolate the temporal trajectories of such events. Second, due to our primary interest to investigate the effects of longer-term duration of abstinence on MDD, we did not include participants with equal or less than 10 years abstinence in this study. Future work may benefit by exploring smokers with differing levels of time abstinent from smoking and its relation to MDD and vice versa. Third, only MDD was studied based on the a priori theoretical model. Thus, it is not possible to generalize the findings from these data to all mood disorders. Accordingly, further evaluation of the associations between different smoking trajectories and other mood (e.g., bipolar) and even anxiety disorders would be advisable. Fourth, we did not have data on MDD severity among participants. Therefore, it is unclear how severity of illness relates to smoking cessation. Fifth, the MIDUS is a survey among persons aged 25–74. It is therefore not clear whether the findings are applicable to other age groups. Sixth, levels of nicotine dependence were not measured in this study. In light of evidence for the differential impact of levels of smoking heaviness [10,58] and nicotine dependence [3,27,59,60] in terms of MDD, future investigations could benefit from examining differential MDD-related predictive validity across levels of smoking heaviness (e.g., heavy smoking, light smoking) [58]. Finally, it is possible that the observed associations could result from unmeasured confounding effects of other MDD-relevant lifestyle factors such as diet, sleep, or exercise [61]. Future examination of other health behaviors in models of smoking-MDD comorbidity would be useful.

## 5. Conclusion

The present findings highlight the clinically important role of smoking in the persistence of MDD over a 10-year period. These results suggest that there may be merit in targeting smoking to reduce the risk of MDD and the mental health benefits of quitting smoking in the form of reduced risk of MDD could usefully be added to common information listed as reasons to quit.

### Declaration of interest

All authors declare that they have no conflicts of interest.

### Acknowledgment

The MIDUS Study was funded by United States Department of Health and Human Services, the National Institutes of Health and the National Institute on Aging. We thank Bianca Jiang for her assistance with manuscript preparation. Work on this study was supported by 2DA20892-A1-06.

## References

- [1] Grant BF, Hasin DS, Chou SP, Stinson FS, Dawson DA. Nicotine dependence and psychiatric disorders in the United States. *Arch Gen Psychiatry* 2004;61:1107-15.
- [2] Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and mental illness: a population-based prevalence study. *JAMA* 2000;284:2606-10.
- [3] Breslau N, Kilbey MM, Andreski P. Nicotine dependence, major depression, and anxiety in young adults. *Arch Gen Psychiatry* 1991;48:1069-74.
- [4] Breslau N, Johnson EO. Predicting smoking cessation and major depression in nicotine-dependent smokers. *Am J Public Health* 2000;90:1122-7.
- [5] Breslau N, Peterson EL, Schultz LR, Chilcoat HD, Andreski P. Major depression and stages of smoking: a longitudinal investigation. *Arch Gen Psychiatry* 1998;55:161-6.
- [6] Rohde P, Kahler CW, Lewinsohn PM, Brown RA. Psychiatric disorders, familial factors, and cigarette smoking: II. Associations with progression to daily smoking. *Nicotine Tob Res* 2004;6:119-32.
- [7] Markou A, Kenny PJ. Neuroadaptations to chronic exposure to drugs of abuse: relevance to depressive symptomatology seen across psychiatric diagnostic categories. *Neurotox Res* 2002;4:297-313.
- [8] Markou A, Kosten TR, Koob GF. Neurobiological similarities in depression and drug dependence: a self-medication hypothesis. *Neuropsychopharmacology* 1998;18:135-74.
- [9] Choi WS, Patten CA, Gillin JC, Kaplan RM, Pierce JP. Cigarette smoking predicts development of depressive symptoms among U.S. adolescents. *Ann Behav Med* 1997;19:42-50.
- [10] Klungsoyr O, Nygard JF, Sorensen T, Sandanger I. Cigarette smoking and incidence of first depressive episode: an 11-year, population-based follow-up study. *Am J Epidemiol* 2006;163:421-32.
- [11] Steuber TL, Banner F. Adolescent smoking and depression: which comes first? *Addict Behav* 2006;31:133-6.
- [12] Covey LS. Comments on 'History of depression and smoking cessation outcome: a meta-analysis'. *Nicotine Tob Res* 2004;6:743-5.
- [13] Hall SM, Humfleet GL, Reus VI, Muñoz RF, Cullen J. Extended nortriptyline and psychological treatment for cigarette smoking. *Am J Psychiatry* 2004;161:2100-7.
- [14] Hitsman B, Borrelli B, McChargue DE, Spring B, Niaura R. History of depression and smoking cessation outcome: a meta-analysis. *J Consult Clin Psychol* 2003;71:657-63.
- [15] Haas AL, Muñoz RF, Humfleet GL, Reus VI, Hall SM. Influences of mood, depression history, and treatment modality on outcomes in smoking cessation. *J Consult Clin Psychol* 2004;72:563-70.
- [16] Cinciripini PM, Wetter DW, Fouladi RT, Blalock JA, Cinciripini LG, Baile WF. The effects of depressed mood on smoking cessation: mediation by postcessation self-efficacy. *J Consult Clin Psychol* 2003;71:292-301.
- [17] Niaura R, Abrams DB. Stopping smoking: a hazard for people with a history of major depression? *Lancet* 2001;357:1900-1.
- [18] MacPherson L, Tull MT, Matusiewicz AK, Rodman S, Strong DR, Kahler CW, et al. Randomized controlled trial of behavioral activation smoking cessation treatment for smokers with elevated depressive symptoms. *J Consult Clin Psychol* 2010;78:55-61.
- [19] Hall SM, Reus VI, Muñoz RF, Sees KL, Humfleet G, Hartz DT, et al. Nortriptyline and cognitive-behavioral therapy in the treatment of cigarette smoking. *Arch Gen Psychiatry* 1998;55:683-90.
- [20] Brown RA, Kahler CW, Zvolensky MJ, Lejuez CW, Ramsey SE. Anxiety sensitivity: relationship to negative affect smoking and smoking cessation in smokers with past major depressive disorder. *Addict Behav* 2001;26:887-99.
- [21] Berlin I, Chen H, Covey LS. Depressive mood, suicide ideation and anxiety in smokers who do and smokers who do not manage to stop smoking after a target quit day. *Addiction* 2010;105:2209-16.
- [22] Wiesbeck GA, Kuhl HC, Yaldizli Ö, Wurst FM. Tobacco smoking and depression—results from the WHO/ISBRA study. *Neuropsychobiology* 2008;57:26-31.
- [23] Sachs-Ericsson N, Collins N, Schmidt B, Zvolensky MJ. Older adults and smoking: characteristics, nicotine dependence and prevalence of *DSM-IV* 12-month disorders. *Aging Ment Health* 2011;15:132-41.
- [24] Sachs-Ericsson N, Schmidt NB, Zvolensky MJ, Mitchell M, Collins N, Blazer DG. Smoking cessation behavior in older adults by race and gender: the role of health problems and psychological distress. *Nicotine Tob Res* 2009;11:433-43.
- [25] Hatsukami DK, Stead LF, Gupta PC. Tobacco addiction. *Lancet* 2008;371:2027-38.
- [26] Korhonen T, Broms U, Varjonen J, Romanov K, Koskenvuo M, Kinnunen T, et al. Smoking behaviour as a predictor of depression among Finnish men and women: a prospective cohort study of adult twins. *Psychol Med J Res Psychiatry Allied Sci* 2007;37:705-15.
- [27] Husky MM, Mazure CM, Paliwal P, McKee SA. Gender differences in the comorbidity of smoking behavior and major depression. *Drug Alcohol Depend* 2008;93:176-9.
- [28] Laaksonen M, Rahkonen O, Karvonen S, Lahelma E. Socioeconomic status and smoking: analysing inequalities with multiple indicators. *Eur J Public Health* 2005;15:262-9.
- [29] Businelle MS, Kendzor DE, Reitzel LR, Costello TJ, Cofta-Woerpel L, Li Y, et al. Mechanisms linking socioeconomic status to smoking cessation: a structural equation modeling approach. *Health Psychol* 2010;29:262-73.
- [30] Holm KE, LaChance HR, Bowler RP, Make BJ, Wamboldt FS. Family factors are associated with psychological distress and smoking status in chronic obstructive pulmonary disease. *Gen Hosp Psychiatry* 2010;32:492-8.
- [31] Prochaska JJ, Delucchi K, Hall SM. A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *J Consult Clin Psychol* 2004;72:1144-56.
- [32] Leeman RF, McKee SA, Toll BA, Krishnan-Sarin S, Cooney JL, Makuch RW, et al. Risk factors for treatment failure in smokers: relationship to alcohol use and to lifetime history of an alcohol use disorder. *Eur J Public Health* 2008;10:1793-809.
- [33] Brim OG, Baltes PB, Bumpass LL, Cleary PD, Featherman DL, Hazzard WR, et al. National survey of midlife development in the United States (MIDUS). Boston, MA: Harvard Medical School, Department of Health Care Policy; 1995–1996.
- [34] Ryff CD, Keyes CLM, Hughes DL. Psychological well-being in MIDUS: profiles of ethnic/racial diversity and life-course uniformity. In: Brim OG, Ryff CD, & Kessler RC, editors. *How healthy are we? A national study of well-being at midlife*. Chicago, IL US: University of Chicago Press; 2004. p. 398-422.
- [35] Kessler RC, Andrews G, Mroczek D, Ustun B, Wittchen HU. The World Health Organization Composite International Diagnostic Interview short-form (CIDI-SF). *Int J Methods Psychiatr Res* 1998;7:171-85.
- [36] Hedden S, Gfroerer J, Barker P, Smith S, Pemberton MR, Saavedra LM, et al. Comparison of NSDUH mental health data and methods with other data sources. CBHSQ Data Review. Center for Behavioral Health Statistics and Quality SAAMHSAS; 2012. [Ed].
- [37] Breslau N, Schultz LR, Johnson EO, Peterson EL, Davis GC. Smoking and the Risk of Suicidal Behavior. *Arch Gen Psychiatry* 2005;62:328-34.
- [38] McClave AK, Dube SR, Strine TW, Kroenke K, Caraballo RS, Mokdad AH. Associations between smoking cessation and anxiety and depression among U.S. adults. *Addict Behav* 2009;34:491-7.
- [39] Khaled SM, Bulloch AG, Williams JV, Hill JC, Lavorato DH, Patten SB. Persistent heavy smoking as risk factor for major depression (MD) incidence—evidence from a longitudinal Canadian cohort of the National Population Health Survey. *J Psychiatr Res* 2012;46:436-43.

- [40] Kahler CW, Spillane NS, Busch AM, Leventhal AM. Time-varying smoking abstinence predicts lower depressive symptoms following smoking cessation treatment. *Eur J Public Health* 2011;13:146-50.
- [41] Fava GA, Rafanelli C, Grandi S, Conti S, Belluardo P. Prevention of recurrent depression with cognitive behavioral therapy: preliminary findings. *Arch Gen Psychiatry* 1998;55:816-20.
- [42] Fava GA, Tomba E, Grandi S. The road to recovery from depression—don't drive today with yesterday's map. *Psychother Psychosom* 2007;76:260-5.
- [43] Ruini C, Fava GA. Clinical applications of well-being therapy. In: Linley PA, & Joseph S, editors. *Positive psychology in practice*. Hoboken, NJ US: John Wiley & Sons Inc.; 2004. p. 371-87.
- [44] Mykletun A, Overland S, Aarø LE, Liabø HM, Stewart R. Smoking in relation to anxiety and depression: evidence from a large population survey: the HUNT study. *Eur Psychiatry* 2008;23:77-84.
- [45] Hämäläinen J, Kaprio J, Isometsä E, Keikkinen M, Poikolainen K, Linderman S, et al. Cigarette smoking, alcohol intoxication and major depressive episode in a representative population sample. *J Epidemiol Community Health* 2001;55:573-6.
- [46] Ford DE, Erlinger TP. Depression and C-reactive protein in US adults: data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2004;164:1010-4.
- [47] Malone KM, Waternaux C, Haas GL, Cooper TB, Li S, Mann JJ. Cigarette smoking, suicidal behavior, and serotonin function in major psychiatric disorders. *Am J Psychiatry* 2003;160:773-9.
- [48] Degenhardt L, Hall W, Lynskey M. Alcohol, cannabis and tobacco use among Australians: a comparison of their associations with other drug use and use disorders, affective and anxiety disorders, and psychosis. *Addiction* 2001;96:1603-14.
- [49] Massak A, Graham K. Is the smoking-depression relationship confounded by alcohol consumption? An analysis by gender. *Eur J Public Health* 2008;10:1231-43.
- [50] Perez-Stable EJ, Marin G, Marin BV, Katz MH. Depressive symptoms and cigarette smoking among Latinos in San Francisco. *Am J Public Health* 1990;80:1500-2.
- [51] Covey LS, Hughes DC, Glassman AH, Blazer DG, George LK. Ever-smoking, quitting, and psychiatric disorders: evidence from the Durham, North Carolina, Epidemiologic Catchment Area. *Tob Control* 1994;3:222-7.
- [52] Kinnunen T, Haukkala A, Korhonen T, Quiles ZN, Spiro A, Garvey AJ. Depression and smoking across 25 years of the normative aging study. *Int J Psychiatry Med* 2006;36:413-26.
- [53] Flensburg-Madsen T, von Scholten MB, Flachs EM, Mortensen EL, Prescott E, Tolstrup JS. Tobacco smoking as a risk factor for depression. A 26-year population-based follow-up study. *J Psychiatr Res* 2011;45:143-9.
- [54] Vandenberg DJ, Bennett CJ, Grant MD, Strasser AA, O'Connor R, Stauffer RL, et al. Smoking status and the human dopamine transporter variable number of tandem repeats (VNTR) polymorphism: failure to replicate and finding that never-smokers may be different. *Eur J Public Health* 2002;4:333-40.
- [55] Munafò MR, Clark TG, Johnstone EC, Murphy MF, Walton RT. The genetic basis for smoking behavior: a systematic review and meta-analysis. *Eur J Public Health* 2004;6:583-97.
- [56] Whitfield JB, Pang D, Bucholz KK, Madden PAF, Heath AC, Statham DJ, et al. Monoamine oxidase: associations with alcohol dependence, smoking and other measures of psychopathology. *Psychol Med J Res Psychiatry Allied Sci* 2000;30:443-54.
- [57] Neuhaus A, Bajbouj M, Kienast T, Kalus P, Von Haebler D, Winterer G, et al. Persistent dysfunctional frontal lobe activation in former smokers. *Psychopharmacology (Berl)* 2006;186:191-200.
- [58] Martini S, Wagner FA, Anthony JC. The association of tobacco smoking and depression in adolescence: evidence from the United States. *Subst Use Misuse* 2002;37:1853-67.
- [59] Breslau N, Kilbey MM, Andreski P. Nicotine dependence and major depression: new evidence from a prospective investigation. *Arch Gen Psychiatry* 1993;50:31-5.
- [60] Breslau N, Kilbey MM, Andreski P. *DSM-III-R* nicotine dependence in young adults: prevalence, correlates and associated psychiatric disorders. *Addiction* 1994;89:743-54.
- [61] Lopresti AL, Hood SD, Drummond PD. A review of lifestyle factors that contribute to important pathways associated with major depression: diet, sleep and exercise. *J Affect Disord* 2013;148:12-27.